

# Bioaccumulation of polychlorinated biphenyls in the eel (*Anguilla anguilla*) at the Camargue Nature Reserve – France

C.A. Oliveira Ribeiro <sup>a,\*</sup>, Y. Vollaire <sup>b</sup>, E. Coulet <sup>c</sup>, H. Roche <sup>b</sup>

<sup>a</sup> Universidade Federal do Paraná, Departamento de Biologia Celular, Caixa Postal 19031, CEP: 81.531-990 Curitiba, PR, Brazil

<sup>b</sup> UMR8079 CNRS-Université Paris-Sud XI, Ecologie Systématique et Evolution, Bât 362, F91405 Orsay Cedex, France

<sup>c</sup> Réserve Nationale de Camargue, La Capelière, F13200 Arles, France

Received 8 March 2007; received in revised form 1 August 2007; accepted 8 August 2007

*The reserve of Camargue – South of France is impacted by a myriad of pollutant organic persistent like PCBs.*

## Abstract

Fish consumption is a potential source of human exposure to pollutants. Here, we study residue levels of PCBs in the eel, *Anguilla anguilla*, from the Nature Camargue Reserve in southern France. Chromatographic analysis (GC-ECD) found seventy identifiable congeners, among which, 10 are considered as dioxin-like PCBs, such as the non-ortho PCB 81 and the mono-ortho chlorobiphenyls PCB105, 114, 118, 123, 156, 157, 167, 170, 180. Toxic Equivalents (TEQ, WHO 2005 TEF-Toxic Equivalent Factors) varied among sites with a maximum in eels from Mornès (29.6 pg g<sup>-1</sup> dry weight). Indicator PCBs (28, 52, 101, 118, 138, 153 and 180) were 22% and 29% of the total PCBs in livers and muscles respectively. Greater homogeneous bioaccumulation in muscle than that in liver suggests an increase risk for humans due to fish consumption.

© 2007 Elsevier Ltd. All rights reserved.

**Keywords:** PCBs; Bioaccumulation; *Anguilla anguilla*; Camargue Reserve; Risk assessments

## 1. Introduction

Polychlorinated Biphenyls (PCBs) are contaminants released to environment due to antropogenic activities and have been reported as members of the group of ubiquitous, persistent, lipophilic, bioaccumulative and toxic or highly toxic microcontaminants (Falandysz et al., 2004). According to Baars et al. (2004) the waste disposal, both of house-holds and industrial wastes, is the major source of PCB emissions into the environment (Baars et al., 2004).

Fish consumption is a potential route of exposure for environmentally persistent organochlorine contaminants and the dietary uptake has contributed significantly to increase the

human health risks (Turyk et al., 2006). In addition the traditional diet of human population in Europe plays an important role in the exposition to a PCB contamination (Johansen et al., 2004), where more than 90% of the total daily intake is derived from food (Papadopoulos et al., 2004). Recently the investigation in aquatic environment has revealed an increase in the bioaccumulation levels of PCBs in fish from different regions of the world (Falandysz et al., 2004; Ashizawa et al., 2005; Borga and Di Gardo, 2005; Evenset et al., 2005; Nakata et al., 2005; Stow et al., 2005; Vives et al., 2005; Ueno et al., 2005; Covaci et al., 2006).

The polychlorinated biphenyls can be accumulated in aquatic organism through the food chain, and its dietary intake by human population is already an evidence as demonstrated in studies with rats (Lehmler et al., 2005), human breast milk (Kunisue et al., 2006), adipose tissue of women (Vaclavik et al., 2006) and human serum levels (Hagmar et al., 2006).

\* Corresponding author. Tel.: +55 41 3361 1680; fax: +55 41 3266 2042.

E-mail address: [ciro@ufpr.br](mailto:ciro@ufpr.br) (C.A. Oliveira Ribeiro).

According to Priha et al. (2005) PCBs have mainly long-term effects and are expected to increase cancer risks, neurodevelopmental and hormonal effects in humans, also including immunosuppression disturbs (Tryphonas et al., 1991). Our previous results (Oliveira Ribeiro et al., 2005) showed a high occurrence of chronic lesions as tumors in liver and spleen and high incidence of lipidosis in liver of *A. anguilla*. Other reported the effects of PCBs as endocrine hormone disrupter (Lin et al., 2006), oxidative stress agent (Murugesan et al., 2005), neurotoxic pollutant (Lehmler et al., 2005), cytochrome P4501A inducer (Behrens and Segner, 2005), a potent birth defects risk (Mendola et al., 2005) and immunomodulator (Iwanowicz et al., 2005).

Due to the complex nature of the 209 congeners the scientific community has been led to classified the representative congeners in two groups: the ‘indicator PCBs’ and the ‘dioxin-like PCBs’. The 7 ‘indicator’ congeners (IUPAC nos 28, 52, 101, 118, 138, 153, 180) have been chosen on the basis of their persistence in food web and their tendency to biomagnification. The 12 dioxin-like PCBs present a common mechanism of action with the polychlorinated dibenzo-*p*-dioxins (PCDDs) and the dibenzofurans (PCDFs): the bind ability to the aryl hydrocarbon (Ah) receptor. Consequently, it is now admitted that the World Health Organization Toxic Equivalency Factors (WHO-TEFs) are the best approach for risk assessment of halogenated aromatic hydrocarbons with dioxin-like properties (Van den Berg et al., 1998, 2005).

The present study will focus the residues levels of PCBs in *A. anguilla* from the Nature Reserve of Camargue – South of France, in order to increase the data about the potential human health risks associated with these contaminants due to fish consumption. This work is part of a large monitoring program supported by INRS and Camargue Nature Reserve and reports the results conducted from October 2003.

## 2. Materials and methods

### 2.1. Studied area and biological material

The Camargue Biosphere Reserve located in the Rhône delta – south of France, occupies 13,000 ha between the northern Vaccarès lagoon and the last undamaged sand dunes of the Mediterranean coast, is the largest coastal wetland area in Western Europe. The studied species was the European eel (*Anguilla anguilla*), a common predator fish living both in contact with sediments and in free waters. Thirty eels (weight  $372 \pm 39$  g; length  $55 \pm 2$  cm) were collected from three riparian locations: Fumemorte canal (FUM), ‘La Capelière’ (CAP) and the Mornès peninsula (MOR), the reference site (Fig. 1) in autumn (October 2003), and organs were carefully excised to chemical analyses.

### 2.2. Polychlorinated biphenyls extraction, separation, analyses procedures and characteristics

Livers and muscles were dissected for PCB congeners analysis and lipid fractions were extracted with a dichloromethane-methanol solution (Folch et al., 1957 modified method). Purification procedures were performed by means of solid phase extraction (SPE) on Florisil column according to the EPA method 3620 (Bond Elut Florisil, 1 g, 200  $\mu$ m particle size – Varian) and analyses were performed by gas chromatography with AutoSystem XL (Perkin-Elmer), using electron capture detection (ECD 63Ni Source) and nitrogen as the carrier gas following an adapted procedure of the EPA Method 8081a. The detection limit ranged from 0.05 to 0.20  $\text{g kg}^{-1}$  in fish tissues.

Among the 209 congeners, 7 compounds are considered as indicators PCBs (PCB 28, 52, 101, 118, 138, 153, 180), representing the major part of the PCB in biological tissues, and 12 compounds as dioxin-like PCB (non-ortho or the mono-ortho PCBs 77, 81, 105, 114, 118, 123, 126, 156, 157, 167, 169, 189 to which are added the PCBs 170 and 180).

### 2.3. TEQ evaluation

The TEQ of dioxin-like PCBs was calculated according Van den Berg et al. (1998) using the WHO-TEFs (World Health Organization – Toxic Equivalent Factors) for fish. The concept of TEF is based the following criteria: (1) show a structural relationship to the PCDDs and PCDFs; (2) bind to the Ah receptor (AhR); (3) elicit AhR mediated biochemical and toxic responses; and (4) be




Fig. 1. The Biophere Camargue National Reserve – South of France: Mor – Mornès site; Cap – La Capelière site; Fum – Fumemorte site.

persistent and accumulate in the food chain. The evaluation of the WHO-TEFs for fish, quite different than those for mammals and birds, give the upper limit value for some PCB ( $<0.00005$ ) (see Table 1), then we have calculated TEFs in excess ( $TEF = 0.00005$ ).

#### 2.4. Statistical procedures

Statistical analyses of PCBs data were achieved with the Statview program (version 5). Due to their non-Gaussian distribution, the data are presented as mean and extreme values. Intersite differences were checked using a non-parametric test, the rank-sum Mann–Whitney  $U$ -test. All tests were regarded as statistically significant when  $p < 0.05$ .

### 3. Results

The distribution of indicator and dioxin-like PCBs in eel muscles and livers from the Camargue Reserve is summarized in Fig. 2 and Table 1. Analysis of lipophilic pollutants in eels from the Vaccarès Lagoon demonstrates clearly inter-site variations of the PCB contamination (Fig. 2, Table 1). Concentration was greater in eels from Mornès and La Capelière than from Fumemorte (Fig. 2). Inter-tissue variation regardless of PCB type is evident in bioaccumulation profiles (indicator, dioxin-like or otherwise), especially in the congeners 19, 52/69, 123, 101/150, 138. Congeners 153 (muscle) and 138 (liver) reached levels  $>100 \text{ ng g}^{-1}$  dry weight (all values are in dry weight). As in muscle, congeners 19 and 138 were most found in livers ( $>90 \text{ ng g}^{-1}$ , Table 1).

PCBs in eels from La Capelière were similar in muscles and livers ( $1300$  and  $1600 \text{ ng g}^{-1}$ , respectively), in contrast with both Mornès and Fumemorte, where bioaccumulation in muscles is greater than in livers ( $2700$  vs.  $1300 \text{ ng g}^{-1}$ , in eels from Mornès;  $1600$  vs.  $900 \text{ ng g}^{-1}$ , in eels from Fumemorte, both  $p < 0.05$ , Table 2). Also, PCB bioaccumulation is more homogeneously distribution in livers than muscles. Individual variability in PCB bioaccumulation is more evident in liver than muscle. These differences seem to be due to indicator PCBs. Indeed, seven indicator congeners were found, often in high concentrations (in order of decreasing concentrations:  $52 > 153 > 138 > 123 > 101 > 180/118$ , Table 1). Indicator PCBs comprised from 4.2% (in liver at Fumemorte) to 45% (muscle from La Capelière) of the total PCBs, with averages of 22% and 29% in livers and muscles respectively, within the entire eel population. Fumemorte typically had eels with lower concentrations (vs. La Capelière and Mornès, both  $p < 0.05$ ; Table 2).

Lower concentrations of  $\sum$ di-CB,  $\sum$ hepta-CB and  $\sum$ octa-CB were found in both tissues from all studied sites, with little or no bioaccumulation, while the congeners  $\sum$ tri-CB,  $\sum$ tetra-CB,  $\sum$ penta-CB, and  $\sum$ hexa-CB were more concentrated (Fig. 3). Bioaccumulation of PCBs was less in eels from Fumemorte than Mornès (liver) or La Capelière (muscle).

Dioxin-like PCBs comprised a small part of total PCBs, from 0.13% in muscle in one individual from Fumemorte and 17% in fish liver from La Capelière, with an average of  $\sim 8\%$ . These dioxin-like congeners were the PCBs 81, 105, 114, 118, 123, 156, 157, 167, 170, 167 and 180. The congeners contributing most to the WHO-TEQ, the PCB 126 and 169

were not detected. Individuals from Mornès had greater concentration of total PCBs in muscle than those from La Capelière ( $p < 0.05$ ). This variation is notably due to the indicator PCBs and the dioxin-like PCBs with greater bioaccumulation levels in muscle at Mornès (Table 2).

WHO-TEQs (1997) values were rather low (Table 2). The greatest mean values were from La Capelière (liver) and Mornès (muscle;  $4.4$  and  $10 \text{ pg g}^{-1}$  dry weight, respectively), while the greatest level was from La Capelière ( $24 \text{ pg g}^{-1}$  dry weight). The recent revised amount of admissible WHO toxic equivalents in foodstuffs is  $4.0 \text{ pg g}^{-1}$  fresh weight in the eel muscle, i.e. about  $12.3 \pm 0.5 \text{ pg g}^{-1}$  dry weight, since muscle contains  $67.5 \pm 1.1$  per cent water. In our study, only two eels from each, Mornès and La Capelière had TEQ levels above that limit (Van den Berg et al., 2005, Table 1).

### 4. Discussion

Polychlorinated biphenyls (PCBs) are aromatic synthetic chemicals, yet they have become common in the biosphere of many parts of the world. While no evidence yet links human death to acute exposure to PCBs, long term occupational exposure can increase mortality from cardiovascular disease and cancer (Pavuk et al., 2004). PCBs are lipophilic compounds, with a  $K_{ow}$  from 4.69–8.05 following their chlorination level (Zhou et al., 2005). Thus, accumulation patterns of each congener is expected to vary since their chemical and physical properties vary. In general, fish and shellfish consumption has contributed significantly to human exposure to these compounds, but more information is necessary to better understand patterns of effects due to different PCB congeners. Therefore, wildlife may be regarded as indicators for human health, providing evidence useful in a human health risk assessment and in identifying new sources of potential cancer-causing agents.

The potential effects of PCBs to human and wildlife is suggested by the serum reaction to adipose tissues that is reported for various PCB congeners (74, 99, 118, 138, 146, 153, 156, 167, 170, 180, 183 and 187). This suggests that humans, aquatic mammals, birds, fish and other biota have a similar PCB profile (ATSDR, 2000). Here, all congeners cited, except CB99, were found in *A. anguilla* from Camargue Reserve, and so fish consumption can be an important source of human exposure to PCBs. From 12 to 14 PCBs are regarded as being particularly toxic because of their ability to bind the aryl hydrocarbon receptor (AhR), a cytosolic receptor protein in most vertebrate tissues. Notably, AhR has a high affinity for 2, 3, 7, 8-substituted PCDD/Fs and some non-ortho and mono-ortho-substituted PCBs. They are called dioxin-like PCBs ranked as a function of the Toxic Equivalent Dioxin (TEQ, Van den Berg et al., 1998, 2005) and include 126, 169, 114, 156, 157, 77, 81, 105, 118, 123, 189, 167 plus 170 and 180. Neither PCB126 nor 169 were found in this study where the majority of eels had low TEQs. Still, risk evaluation based on TEQ dioxin is still debated since some (e.g., PCB126) have low response levels for enzyme induction in human cells, including primary hepatocytes, breast cancer cell lines and primary lymphocytes (Zeiger et al., 2001).

Table 1  
Summary of PCBs compounds found in liver and muscle of *Anguilla anguilla* collected in Camargue Reserve – South of France, at October, 2003

Tissue site	1997 WHO-TEFs for fish*	Liver			Muscle			
		La Capelière (8)	Fumemorte (10)	Mornès (10)	La Capelière (11)	Fumemorte (9)	Mornès (8)	
CB8		15.0 [1.1–63.4]	14.6 [2.5–31.3]	20.3 [9.2–59.6]	CB8	14.7 [3.3–43.7]	27.9 [2.2–65.5]	18.8 [6.9–46.9]
CB14		20.8 [2.3–75.2]	75.4 [nd–387]	26.5 [nd–127]	CB14	8.2 [nd–48.0]	9.2 [3.3–26.5]	8.9 [0.0–20.2]
CB13/15		11.4 [nd–21.0]	71.4 [8.6–306]	29.3 [2.2–81.4]	CB13/15 <sup>a,b</sup>	4.3 [0.2–16.3]	28.9 [15.5–65.3]	27.6 [2.4–69.1]
CB19 <sup>b,c</sup>		56.8 [20.7–94.5]	81.1 [21.2–133]	27.1 [nd–70.4]	CB19 <sup>a,b</sup>	48.4 [0.8–128]	130.4 [nd–240]	252.2 [11.7–596]
CB18		9.2 [nd–24.1]	4.4 [nd–13.4]	6.0 [nd–15.8]	CB18 <sup>a</sup>	2.8 [nd–10.4]	32.3 [1.9–101.7]	10.0 [nd–48.4]
CB17		11.9 [nd–27.0]	26.0 [nd–92.0]	15.9 [nd–49.7]	CB17	7.8 [nd–22.7]	18.7 [nd–48.8]	15.4 [nd–42.0]
CB32		3.0 [nd–6.1]	2.1 [nd–11.6]	2.5 [nd–6.2]	CB32 <sup>a,b</sup>	2.5 [nd–5.9]	7.8 [4.2–19.4]	12.9 [0.6–30.6]
CB16		20.8 [nd–115]	3.2 [nd–9.1]	16.6 [nd–60.1]	CB16 <sup>a</sup>	7.7 [nd–35.3]	8.3 [2.6–20.3]	6.5 [nd–19.2]
CB26/50 <sup>a,c</sup>		20.5 [2.2–54.2]	103.0 [11.3–311]	30.3 [nd–99.9]	CB26/50 <sup>b</sup>	20.4 [nd–51.8]	37.7 [9.8–87.5]	81.0 [13.9–141]
CB25		60.7 [2.4–184.8]	60.6 [3.7–213.0]	86.1 [3.6–410]	CB25	14.6 [nd–71.4]	14.0 [3.1–36.8]	13.2 [1.8–33.3]
CB51		5.3 [nd–17.9]	5.5 [nd–35.6]	5.5 [nd–34.2]	CB51	8.9 [nd–27.7]	3.8 [nd–19.8]	1.7 [nd–10.1]
CB31 <sup>c</sup>		24.7 [nd–173]	0.8 [nd–8.2]	9.0 [nd–32.2]	CB31	13.7 [nd–30.7]	14.2 [3.6–65.8]	10.4 [2.0–27.5]
CB28 <sup>d</sup>		76.2 [4.9–274]	13.1 [1.9–37.2]	17.3 [nd–46.9]	CB28	63.0 [nd–202]	68.8 [19.3–296]	36.7 [6.1–149]
CB36		17.6 [nd–59.7]	9.3 [1.8–22.4]	26.6 [nd–98.6]	CB36	10.0 [2.1–22.8]	9.6 [4.0–21.0]	16.4 [2.0–34.2]
CB20/33/45 <sup>c</sup>		9.2 [nd–23.5]	4.8 [1.6–10.9]	10.4 [0.9–19.1]	CB20/33/45	7.8 [0.3–19.5]	11.9 [3.8–49.8]	5.9 [1.2–12.1]
CB22 <sup>c</sup>		9.8 [nd–34.3]	22.2 [5.2–71.4]	17.0 [0.7–71.2]	CB22 <sup>a</sup>	12.0 [3.3–33.4]	2.6 [nd–7.1]	13.5 [5.9–26.9]
CB52/69 <sup>c</sup>		25.0 [12.4–60.6]	35.7 [6.4–214]	34.6 [17.0–60.6]	CB52/69	83.1 [13.9–443]	37.8 [21.6–116]	229.1 [13.8–843]
CB43/49		11.3 [2.2–17.7]	7.5 [nd–18.8]	14.0 [2.7–30.7]	CB43/49	12.7 [1.7–27.4]	10.0 [3.9–22.2]	5.6 [nd–15.0]
CB47/48/65 <sup>c</sup>		4.0 [nd–10.5]	1.5 [nd–3.5]	6.1 [1.4–13.7]	CB47/48/65 <sup>c</sup>	9.1 [nd–37.6]	7.5 [2.7–22.8]	2.4 [nd–8.4]
CB44 <sup>a,c</sup>		8.8 [0.6–28.6]	2.0 [nd–5.3]	6.3 [1.2–12.7]	CB44	10.5 [nd–36.5]	34.6 [4.0–201]	12.1 [nd–18.5]
CB42		5.2 [nd–24.8]	2.7 [nd–6.6]	3.1 [nd–14.1]	CB42	4.0 [nd–17.5]	10.6 [0.5–34.8]	4.3 [0.14–10.0]
CB64/72		8.5 [2.0–17.8]	8.3 [1.4–20.7]	13.3 [1.1–31.6]	CB64/72 <sup>a</sup>	9.1 [nd–28.9]	17.4 [11.1–27.0]	12.7 [4.5–20.0]
CB71/103		5.9 [nd–22.4]	3.3 [nd–9.3]	7.9 [nd–18.8]	CB71/103 <sup>c</sup>	22.9 [nd–92.3]	13.6 [1.8–53.8]	4.3 [nd–19.7]
CB37/41/68		13.5 [nd–99.6]	0.9 [nd–4.1]	0.4 [nd–1.2]	CB37/41/68	3.1 [nd–18.3]	0.6 [nd–1.9]	1.7 [nd–5.9]
CB40		0.5 [nd–2.6]	0.2 [nd–2.1]	0.4 [nd–2.2]	CB40 <sup>a,c</sup>	0.02 [nd–0.15]	2.0 [nd–6.7]	nd
CB95		45.1 [12.8–122]	18.8 [nd–43.7]	68.1 [nd–382]	CB95 <sup>b,c</sup>	5.5 [nd–17.8]	1.7 [nd–5.9]	27.0 [1.0–92.8]
CB74		59.1 [6.1–218]	33.6 [nd–64.3]	38.1 [4.5–83.1]	CB74	37.8 [2.9–89.2]	58.6 [16.9–168]	60.5 [nd–138]
CB70 <sup>c</sup>		61.4 [12.3–18]	20.4 [7.3–34.9]	45.5 [4.7–84.4]	CB70	52.8 [2.2–139.3]	62.4 [8.0–180]	47.2 [12.3–119]
CB76/80 <sup>c</sup>		70.1 [20.4–184]	15.7 [nd–54.2]	60.6 [4.1–136]	CB76/80 <sup>a</sup>	52.4 [4.8–129]	64.9 [nd–211]	90.8 [16.0–199]
CB66 <sup>c</sup>		5.9 [nd–17.7]	7.7 [nd–41.7]	2.5 [nd–14.9]	CB66	12.9 [nd–90.5]	7.6 [nd–21.7]	16.0 [nd–53.9]
CB92 <sup>c</sup>		35.9 [nd–128]	10.8 [nd–25.6]	27.5 [nd–55.0]	CB92	28.9 [1.4–74.5]	37.4 [nd–140]	48.5 [6.2–85.2]
CB84		4.1 [nd–16.3]	1.8 [nd–9.4]	3.0 [nd–16.3]	CB84	2.9 [nd–6.8]	10.8 [nd–46.5]	9.5 [nd–22.6]
CB101/150 <sup>d,c</sup>		44.5 [2.1–143]	6.7 [nd–19.5]	35.5 [3.8–100]	CB101/150	43.5 [5.3–89.1]	51.6 [4.5–120]	127.0 [12.1–278]
CB60/113		12.5 [3.3–41.0]	7.3 [nd–14.7]	13.7 [1.9–26.9]	CB60/113	9.2 [0.8–22.9]	24.4 [nd–86.1]	23.5 [2.9–45.0]
CB112/119		14.7 [nd–109.3]	0.5 [nd–3.5]	2.8 [nd–7.9]	CB112/119	1.1 [nd–5.3]	0.2 [nd–1.2]	3.6 [nd–12.2]
CB148		7.3 [nd–36.7]	1.0 [nd–2.9]	1.6 [nd–7.8]	CB148	3.3 [nd–15.5]	0.7 [nd–3.4]	2.5 [0.1–8.4]
CB97/117 <sup>a,c</sup>		9.6 [nd–27.2]	1.8 [nd–4.4]	7.2 [1.1–16.1]	CB97/117	10.2 [1.2–48.0]	6.5 [nd–16.4]	37.8 [1.6–101.4]
CB87/115 <sup>a,c</sup>		11 n,0.1 [1.1–42.6]	2.3 [nd–4.5]	9.8 [0.8–26.2]	CB87/115 <sup>b</sup>	11.3 [1.0–24.4]	14.9 [nd–43.0]	32.6 [3.2–60.5]
CB85		10.5 [nd–42.6]	3.1 [nd–13.0]	1.2 [nd–3.1]	CB85 <sup>b</sup>	4.8 [nd–13.4]	15.8 [nd–46.9]	20.4 [0.3–72.9]
CB81 <sup>a,c</sup>	0.0005	7.2 [nd–24.7]	0.1 [nd–1.1]	2.8 [nd–15.8]	CB81 <sup>a,c</sup>	12.7 [1.0–45.9]	4.2 [nd–22.6]	18.0 [2.4–36.1]
CB110 <sup>a,c</sup>		33.0 [7.9–90.4]	7.7 [2.6–20.5]	26.0 [3.1–60.4]	CB110 <sup>b</sup>	28.5 [3.3–59.6]	27.6 [9.3–57.8]	75.6 [9.3–139]
CB151 <sup>a,c</sup>		24.7 [nd–94.2]	2.5 [nd–10.7]	24.2 [2.1–76.8]	CB151 <sup>b</sup>	29.5 [3.1–66.2]	29.6 [nd–62.1]	69.5 [5.3–123]
CB135/144 <sup>a,c</sup>		18.9 [nd–82.8]	2.7 [nd–12.6]	11.8 [1.7–55.7]	CB135/144	20.5 [2.2–46.2]	20.8 [nd–41.3]	53.9 [3.7–95.9]
CB123 <sup>a,c</sup>	< 0.000005	68.3 [14.8–226]	12.5 [1.6–50.3]	55.5 [4.6–159]	CB123 <sup>b</sup>	46.0 [5.3–112]	58.5 [nd–119]	134 [10.2–231]
CB118 <sup>b,c</sup>		48.2 [nd–100.7]	16.8 [nd–38.7]	15.4 [nd–147]	CB118 <sup>a</sup>	1.6 [nd–6.2]	0.4 [nd–4.0]	0.5 [nd–3.7]
CB146 <sup>b</sup>		37.7 [nd–248]	11.4 [nd–63.7]	48.7 [nd–85.9]	CB146 <sup>c</sup>	44.7 [nd–127]	76.6 [nd–253]	99.3 [6.1–257]
CB114 <sup>b</sup>	< 0.000005	4.4 [nd–12.3]	2.4 [nd–10.6]	4.6 [nd–14.6]	CB114 <sup>b</sup>	0.6 [0.03–2.2]	6.0 [nd–24.3]	5.9 [0.2–17.2]
CB132/179		13.4 [nd–78.4]	4.3 [nd–14.0]	13.0 [0.08–37.0]	CB132/179 <sup>c</sup>	28.9 [1.8–75.8]	23.0 [nd–58.2]	41.3 [nd–92.7]
CB153 <sup>c</sup>		67.3 [0.3–370.0]	14.4 [1.2–70.9]	62.0 [2.5–254]	CB153 <sup>a,b,c</sup>	69.2 [6.0–165]	14.0 [nd–73.2]	193.8 [22.1–369]
CB141/176 <sup>b</sup>		12.1 [nd–47.6]	2.6 [nd–23.4]	48.1 [nd–306]	CB141/176	13.9 [nd–47.1]	3.1 [nd–10.2]	23.1 [nd–96.2]

(continued on next page)



Table 1 (continued)

Tissue site	1997 WHO-TEFs for fish*		Liver				Muscle			
	La Capelière (8)	Fumemorte (10)	Mornès (10)	La Capelière (11)	Fumemorte (9)	Mornès (8)				
<b>CB105<sup>c</sup></b>	<b>26.5 [1.4–107]</b>	<b>7.7 [nd–17.2]</b>	<b>21.3 [1.6–58.2]</b>	<b>26.9 [1.9–62.8]</b>	<b>37.7 [nd–111.1]</b>	<b>38.8 [7.4–119]</b>				
CB137/141 <sup>b,c</sup>	43.4 [nd–176]	7.2 [nd–26.7]	26.6 [3.5–112]	30.8 [1.6–86.0]	51.2 [nd–96.2]	77.5 [5.3–136]				
CB130/178	2.8 [nd–18.6]	0.1 [nd–1.5]	2.2 [nd–21.4]	2.8 [nd–18.3]	0.1 [nd–0.7]	8.1 [nd–19.5]				
CB163/164	2.7 [nd–8.6]	5.6 [nd–21.9]	2.9 [nd–16.8]	1.8 [0.4–6.1]	0.5 [nd–1.9]	8.2 [1.1–25.1]				
<b>CB138<sup>c</sup></b>	<b>105 [nd–497]</b>	<b>21.0 [nd–110]</b>	<b>73.4 [7.3–273]</b>	<b>101 [3.2–246]</b>	<b>64.2 [nd–169]</b>	<b>180.3 [22.9–392]</b>				
CB175/182/187 <sup>b,c</sup>	17.4 [nd–66.1]	2.5 [nd–21.9]	11.4 [nd–58.8]	1.2 [nd–5.0]	13.6 [nd–114.8]	26.6 [1.2–55.0]				
CB166/183 <sup>b,c</sup>	6.8 [nd–36.5]	0.9 [nd–8.7]	3.0 [nd–22.0]	0.5 [nd–2.1]	7.9 [nd–30.0]	20.4 [2.2–33.9]				
CB159/185	0.5 [nd–2.8]	0.1 [nd–1.0]	0.1 [nd–0.4]	1.1 [nd–6.0]	15.9 [nd–69.8]	5.0 [nd–23.3]				
CB128/162/174 <sup>c</sup>	20.8 [nd–98.1]	5.1 [nd–32.0]	20.0 [1.0–60.9]	28.3 [1.5–131]	30.1 [nd–76.5]	56.0 [6.5–107]				
CB177 <sup>c</sup>	18.5 [nd–95.4]	2.8 [nd–21.6]	7.9 [0.7–35.6]	8.4 [nd–25.6]	31.3 [nd–131]	59.2 [1.6–130]				
CB197/201	5.4 [nd–13.5]	1.9 [nd–14.4]	2.2 [nd–10.9]	8.6 [0.02–31.5]	0.7 [nd–1.6]	8.6 [0.7–20.0]				
<b>CB167<sup>b</sup></b>	<b>0.7 [nd–3.6]</b>	<b>nd</b>	<b>0.3 [nd–3.2]</b>	<b>0.1 [nd–0.4]</b>	<b>0.1 [nd–1.2]</b>	<b>0.1 [nd–0.8]</b>				
<b>CB156/157</b>	<b>10.0 [nd–77.4]</b>	<b>0.6 [nd–5.8]</b>	<b>1.3 [nd–7.1]</b>	<b>7.5 [nd–65.0]</b>	<b>0.7 [nd–1.6]</b>	<b>16.6 [0.2–62.6]</b>				
<b>CB180</b>	<b>26.5 [nd–146.2]</b>	<b>4.4 [nd–40.9]</b>	<b>10.5 [nd–45.4]</b>	<b>13.4 [0.8–53.8]</b>	<b>29.7 [nd–72.9]</b>	<b>48.5 [0.8–95.1]</b>				
CB199 <sup>b</sup>	18.1 [nd–90.8]	nd	3.5 [nd–30.7]	3.4 [nd–20.5]	4.4 [nd–21.8]	25.0 [0.4–69.0]				
<b>CB170<sup>c</sup></b>	<b>5.7 [nd–19.7]</b>	<b>1.5 [nd–14.6]</b>	<b>1.1 [nd–5.4]</b>	<b>7.9 [nd–34.7]</b>	<b>0.8 [nd–5.4]</b>	<b>5.9 [nd–16.5]</b>				
CB190/203	2.8 [nd–9.9]	0.5 [nd–5.4]	2.5 [nd–8.6]	0.5 [nd–2.1]	0.7 [nd–2.5]	4.1 [0.2–10.0]				
CB195	8.3 [nd–58.7]	0.9 [nd–4.6]	6.4 [nd–59.7]	0.9 [nd–7.2]	0.6 [nd–1.5]	16.1 [nd–118]				
CB194	51.7 [nd–266]	23.5 [nd–71.2]	21.3 [2.7–58.8]	30.3 [0.9–121.8]	82.1 [9.0–208]	53.9 [4.2–147]				
CB205	0.1 [nd–1.2]	nd	1.3 [nd–13.2]	nd	nd	nd				

Data are expressed as mean and extreme values [min–max]. Conclusions of the Mann–Whitney *U*-test: significant intersite variations  $p < 0.05$ . a: La Capelière vs Fumemorte; b: La Capelière vs Mornès; c: Fumemorte vs Mornès. Italic characters: Indicator PCBs; bold characters: dioxin-like PCBs. \* after van den Berg et al. (1998) the value 1 is attributed to 2,3,7,8-TCDD (tetrachlorodibenzodioxin).

Several dioxin-like compounds on oestrogen metabolism in the malignant MCF 7 and non-tumorigenic MCF 10, a human mammary epithelial cell lines, have been studied (Van Duursen et al., 2003).

The majority of PCBs congeners in eels from the Camargue Nature Reserve are compounds that are still lacking information on toxic potential for humans and wildlife. The greatest PCB concentrations (liver from La Capelière, muscle from Mornès, 4.5  $\mu\text{g g}^{-1}$  dry weight) are greater than both, the AEL (Adverse Effect Level) of 0.007  $\mu\text{g g}^{-1} \text{day}^{-1}$  responsible for neurobehavioral alterations in infant monkeys, and the LOAEL (Lowest Observed Adverse Effect Level) of 0.005  $\mu\text{g g}^{-1} \text{day}^{-1}$  producing immunological effects in adults (ATSDR, Agency for Toxic Substances and Disease Registry, 2000).

PCB contamination varies widely from site to site. Fish with lower concentrations (indicator and dioxin-like PCBs) were sampled near the Fumemorte catchment which receives Rhone River irrigation waters from nearby agriculture. In addition, hepatic and muscular TEQ are also lower than from La Capelière or Mornès. Indeed, fish captured at Mornès, which should not have agriculture or urban water input, had bioaccumulated more PCBs, some of which were dioxin-like (for example 81, 123, 114, 167, 156/157). This unexpected contamination may be due to atmospheric transfer (Banas et al., 2005), to water currents linked to a complex hydrological balance (Chauvelon, 1998), or to biomagnification as top-predators (Iannuzzi et al., 1996; Persic, 2004).

In addition to dioxin-like congeners, the seven indicator PCBs (138, 153 and 28, 52, 101, 118, 180) provide a contamination image. According to WHO (2000), the PCBs 138 and 153 are the major congeners found in all animal samples (as here, where PCB 153 was in both liver and muscle). These cause histological damage in organs including the liver and are neurological and neurochemical toxic agents (Sánchez-Alonso et al., 2003; Duffy and Zelikoff, 2006). A high incidence of necrosis in liver, along with other, organochlorine pesticide contaminants, can occur due to these congeners (Oliveira Ribeiro et al., 2005). Congener 28, which may also damage organs, was found in liver and muscle in this study, and its bioaccumulation is greatest in muscle, and so implies a human health risk. Also, liver concentrations were above the minimal risk level (0.005  $\mu\text{g g}^{-1}$ ). Although the congeners 118, 138, 153, 156, 170, 180 and 185 had relatively high levels in porpoise carp, bird, oligochaete, shrimp and human fat and milk, congener 74 found here in liver and muscle, was the most abundant in human fat and serum (ATSDR, 2000). PCB 153 in the majority of species is due to its slow rate of biotransformation and elimination (Dip et al., 2003; Hoekstra et al., 2003). Therefore, consumption of fish from Vaccarès lagoon is a potential pathway for human health risk.

Effects in liver of rat, mice and monkeys were observed after exposure to PCB 153 and others (Vezina et al., 2004). Exposure to mixtures of PCBs causes a diversity of effects, including hormonal, immunological, neurological and reproductive system damage. PCBs may cause tumor activity and the environmental exposure has been associated with cancer of adipose tissues (Pavuk et al., 2004).




Fig. 2. Distribution of PCBs congeners in muscle and liver of *Anguilla anguilla* from Camargue Reserve — South of France, collected at October, 2003.

Wildlife response to exposure to PCBs and congeners varies widely, possibly reflecting not only variation in susceptibility but also different mechanisms of action or selective metabolism of individual congeners. PCBs tend to

biomagnify in the food chain (Binelli and Proveni, 2003; Bureau et al., 2004), reaching greater toxicity at higher trophic levels, such as in piscivorous and fatty fish (e.g., *A. anguilla*).

Table 2

Comparative total PCBs, indicator PCBs, dioxin-like PCBs and TEQ values in liver and muscle of *Anguilla anguilla* from Camargue Reserve — South of France collected at October 2003

	$\Sigma$ PCBs ng g <sup>-1</sup> dw	$\Sigma$ indicator PCBs ng g <sup>-1</sup> dw	% indicator PCBs	$\Sigma$ dioxin-like PCBs ng g <sup>-1</sup> dw	% dioxin-like PCBs	TEQ pg g <sup>-1</sup> dw
Liver		a,c*	a	a	a,b	a,c*
Vaccarès lagoon (28)	1499 [144–4539]	293 [43–1842]	22.1 [4.2–41]	110.1 [9–646]	8.3 [1.8–17]	2.02 [0.04–13]
La Capelière (8)	1556 [352–4539]	497 [68–1842]	28.1 [7–41]	190.3 [37–646]	11.6 [7–17]	4.4 [0.1–13]
Fumemorte (10)	855 [244–1912]	133 [43–375]	16.6 [4.2–30]	45.9 [9–139]	6.4 [1.8–11]	0.26 [0.04–1]
Mornès (10)	1281 [144–3005]	289 [51–1149]	22.7 [9–38]	110 [9–406]	7.6 [3.7–14]	1.89 [0.04–10]
Muscle	b	c*	a, c	b,c*		c
$\Sigma$ Vaccarès lagoon (27)	1803 [254–4477]	588 [54–1819]	29.3 [10–45]	158 [0.3–468]	8.5 [0.13–14]	6.5 [0.3–24]
La Capelière (10)	1290 [254–2680]	494 [54–1200]	32.9 [15–45]	107 [11–233]	8.6 [4.5–13]	7.1 [0.6–24]
Fumemorte (9)	1554 [774–2997]	331 [89–816]	21.0 [10–29]	134 [0.3–250]	7.9 [0.13–14]	2.6 [0.3–12]
Mornès (8)	2724 [515–4477]	996 [116–1819]	33.9 [20–41]	250 [23–468]	9.1 [4.9–13]	10 [1.3–20]

The data are expressed as mean and extreme values [min–max]. Conclusions of the Mann–Whitney *U*-test: significant intersite variations  $p < 0.05$ . a: La Capelière vs Fumemorte; b: La Capelière vs Mornès; c: Fumemorte vs Mornès. c\*  $p < 0.06$  Fumemorte vs Mornès.




Fig. 3. Comparative distribution of di, tri, tetra, penta, hexa, hepta, octa groups and total PCBs in muscle and liver of *Anguilla anguilla* from La Capelière, Fumemorte and Mornès sites in Camargue Reserve — South of France, collected at October, 2003.

Sport fish consumption in the Great Lakes is therefore a potential source of exposure for environmentally persistent organochlorine contaminants that may cause human health risks (Turyk et al., 2006). Associations occur between fish consumption and body burdens of PCBs within sports or professional fishermen, or others that consume large quantities of contaminated fish (Vaclavik et al., 2006). Advancements in TEQ calculations seem to provide an accurate evaluation of risk for PCBs and dioxin-like molecules in the environment (Alcock et al., 1998).

## 5. Conclusion

PCB exposure from fish consumption is an important source of human exposure and should be better studied. Since certain hydroxylated methylsulfonyl (MeSO<sub>2</sub>) PCB metabolites may reach greater levels than their respective parent compounds, and therefore it is necessary to investigate potential biological or toxicological activities of these persistent metabolites. In addition, the persistence and mobility of these toxic chemicals indicate that the problems are global and that the solution must be at the global level.

## Acknowledgements

This work was supported by CNRS (Centre National de la Recherche Scientifique — France) and CAPES (Brazilian Agency for Science and Technology). We are greatly thankful

to the staff of the National Nature Reserve of Camargue, especially François Lescuyer, for his technical field assistance during fish sampling.

## References

- Alcock, R.E., Behnisch, P.A., Jones, K.C., Hagenmaier, H., 1998. Dioxin like PCBs in the environment: human exposure and the significance of sources. *Chemosphere* 37, 1457–1472.
- Ashizawa, A., Hicks, H.E., De Rosa, C.T., 2005. Human health research and policy development: experience in the Great Lakes region. *International Journal of Hygiene and Environmental Health* 208, 7–13.
- ATSDR, 2000. Toxicological Profile for Polychlorinated Biphenyls (PCBs). US Department of Health and Human Services, Atlanta, USA.
- Baars, A.J., Bakker, M.I., Baumann, R.A., Boon, P.E., Freijer, J.I., Hoogenboom, L.A.P., Hoogerbrugge, R., van Klaveren, J.D., Liem, A.K.D., Traag, W.A., de Vries, J., 2004. Dioxins, dioxin-like PCBs and non-dioxin-like PCBs in foodstuffs: occurrence and dietary intake in The Netherlands. *Toxicology Letters* 151, 51–61.
- Banas, D., Grillas, P., Auby, I., Lescuyer, F., Coulet, E., Moreteau, J.-C., Millet, B., 2005. Short time scale changes in underwater irradiance in a wind-exposed lagoon (Vaccarès lagoon, France): efficiency of infrequent field measurements of water turbidity or weather data to predict irradiance in the water column. *Hydrobiologia* 551, 3–16.
- Behrens, A., Segner, H., 2005. Cytochrome P4501A induction in brown trout exposed to small streams of an urbanized area: results of a five-years-study. *Environmental Pollution* 136, 231–242.
- Binelli, A., Provini, A., 2003. The PCB pollution of Lake Iseo (N. Italy) and the role of biomagnification in the pelagic food web. *Chemosphere* 53, 143–151.
- Borga, K., Di Gardo, A., 2005. Comparing measured and predicted PCB concentrations in Arctic seawater and marine biota. *Science of the Total Environment* 342, 281–300.
- Burreau, S., Zeb, Y., Broman, D., Ishaq, R., 2004. Biomagnification of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) studied in pike (*Esox lucius*), perch (*Perca fluviatilis*) and roach (*Rutilus rutilus*) from the Baltic Sea. *Chemosphere* 55, 1043–1052.
- Chauvelon, P., 1998. A wetland managed for agriculture as an interface between the Rhône river and the Vaccarès lagoon (Camargue, France): transfers of water and nutrients. *Hydrobiologia* 373/374, 181–191.
- Covaci, A., Voorspoels, S., Thomsen, C., van Bavel, B., Neels, G., 2006. Evaluation of total lipids using enzymatic methods for the normalization of persistent organic pollutant levels in serum. *Science of the Total Environment* 366, 361–366.
- Dip, R., Hegglin, D., Deplazes, P., Dafflon, O., Koch, H., Naegeli, H., 2003. Age- and sex-dependent distribution of persistent organochlorine pollutants in urban foxes. *Environmental Health Perspectives* 111, 1608–1612.
- Duffy, J., Zelikoff, J., 2006. The relationship between noncoplanar PCB-induced immunotoxicity and hepatic CYP1A induction in a fish model. *Journal of Immunotoxicology* 3, 39–47.
- Evenset, A., Christensen, G.N., Kallenborn, R., 2005. Selected chlorobornanes, polychlorinated naphthalenes and brominated flame retardants in Bjornoya (Bear Island) freshwater biota. *Environmental Pollution* 136, 419–430.
- Falandysz, J., Wyrzykowska, B., Warzocha, J., Barska, I., Gargacik-Wesolowska, A., Szefer, P., 2004. Organochlorine pesticides and PCBs in perch *Perca fluviatilis* from the Odra/Oder river estuary, Baltic Sea. *Food Chemistry* 87, 17–23.
- Folch, J., Less, M., Sloane-Stanley, G.H., 1957. A simple method for the isolation and purification of total lipids from animal tissues. *Journal Biology Chemistry* 226, 497–509.
- Hagmar, L., Wallin, E., Vessby, B., Jonsson, B.A.G., Bergman, A., Rylander, L., 2006. Intra-individual variations and time trends 1991–2001 in human serum levels of PCB, DDE and hexachlorobenzene. *Chemosphere* 64, 1507–1513.
- Hoekstra, P.F., O'Hara, T.M., Karlsson, H., Solomon, K.R., Muir, D.C.G., 2003. Enantiomer-specific biomagnification of alpha-hexachlorocyclohexane and

- selected chiral chlordane-related compounds within an arctic marine food web. *Environmental Toxicology and Chemistry* 22, 2482–2491.
- Iannuzzi, T.J., Harrington, N.W., Shear, N.M., Curry, C.L., Carlson-Lynch, H., Henning, M.H., Su, S.H., Rabbe, D.E., 1996. Distribution of key exposure factors controlling the uptake of xenobiotic chemicals in an estuarine food web. *Environmental Toxicology and Chemistry* 15, 1979–1992.
- Iwanowicz, L.R., Lerner, D., Blazer, V.S., McCorminck, S.D., 2005. Aqueous exposure to Aroclor 1254 modulates the mitogenic response of Atlantic salmon anterior kidney T-cells: indications of short- and long-term immunomodulation. *Aquatic Toxicology* 72, 305–314.
- Johansen, P., Muir, D., Asmund, G., Riget, F., 2004. Human exposure to contaminants in the traditional Greenland diet. *Science of the Total Environment* 331, 189–206.
- Kunisue, T., Muraoka, M., Ohtake, M., Sudaryano, A., Minh, N.H., Ueno, D., Kigaki, Y., Ochi, M., Tsydenova, O., Kamikawa, S., Tonegi, T., Nakamura, Y., Shimomura, H., Nagayama, J., Tanabe, S., 2006. Contamination status of persistent organochlorines in human breast milk from Japan: recent levels and temporal trend. *Chemosphere* 64, 1601–1608.
- Lehmler, H.-J., Roverton, L.W., Garrison, A.W., Kodavanti, P.R.S., 2005. Effects of PCB 84 enantiomers on [<sup>3</sup>H]-phorvol ester binding in rat cerebellar granule cells and <sup>45</sup>Ca<sup>2+</sup>-uptake in rat cerebellum. *Toxicology Letters* 156, 391–400.
- Lin, Y.-J., Liu, H.-C., Hseu, Z.-Y., Wu, W.-J., 2006. Study of transportation and distribution of PCBs using an ecologically simulated growth chamber. *Chemosphere* 64, 567–573.
- Mendola, P., Robinson, L.K., Buck, G.M., Druschel, C.M., Fitzgerald, E.F., Sever, L.E., Vena, J.E., 2005. Birth defects risk associated with maternal sport fish consumption: potential effect modification by sex of offspring. *Environmental Research* 97, 134–141.
- Murugesan, P., Kanagaraj, P., Yuraj, S., Balasubramanian, K., Aruldas, M.M., Arunakaran, J., 2005. The inhibitory effects of polychlorinated biphenyl Aroclor 1254 on Leydig cell LH receptors, steroidogenic enzymes and antioxidant enzymes in adult rats. *Reproductive Toxicology* 20, 117–126.
- Nakata, H., Hirakawa, Y., Kawazoe, M., Nakabo, T., Arizono, K., Abe, S.-I., Kitano, T., Shimada, H., Watanabe, I., Li, W., Ding, W., 2005. Concentrations and compositions of organochlorine contaminants in sediments, soils, crustaceans, fishes and birds collected from Lake Tai, Hangzhou Bay and Shanghai city region, China. *Environmental Pollution* 133, 415–429.
- Oliveira Ribeiro, C.A., Vollaire, Y., Sanchez-Chardi, A., Roche, H., 2005. Bioaccumulation and the effects of organochlorine pesticides, PAH and heavy metals in the Eel (*Anguilla anguilla*) at the Camargue Nature Reserve, France. *Aquatic Toxicology* 74, 53–69.
- Papadopoulos, A., Vassiliadou, I., Costopoulou, D., Papanicolaou, C., Leondiadis, L., 2004. Levels of dioxins and dioxin-like PCBs in food samples on the Greek market. *Chemosphere* 57, 413–419.
- Pavuk, M., Cerhan, J.R., Lynch, C.F., Schecter, A., Petrik, J., Chovancova, J., Kocan, A., 2004. Environmental exposure to PCBs and cancer incidence in eastern Slovakia. *Chemosphere* 54, 1509–1520.
- Persic, A., 2004. Trophic transfer of organochlorine compounds throughout the aquatic food web of the Camargue Biosphere Reserve – a stable isotope approach. Thesis University of Paris-Sud 11, France.
- Priha, E., Hellman, S., Sorvari, J., 2005. PCB contamination from polysyllphide sealants in residential areas exposure and risk assessment. *Chemosphere* 59, 537–543.
- Sánchez-Alonso, J., López-Aparicio, P., Recio, M., Pérez-Albarsanz, M., 2003. Apoptosis-mediated neurotoxic potential of a planar (PCB 77) and a nonplanar (PCB 153) polychlorinated biphenyl congeners in neuronal cell cultures. *Toxicology Letters* 144, 337–349.
- Stow, J.P., Sova, J., Reimer, K.J., 2005. The relative influence of distant and local (DEW-line) PCB sources in the Canadian Arctic. *Science of the Total Environment* 342, 107–118.
- Tryphonas, H., Luster, M.I., Schiffman, G., Dawson, L.L., Hodgen, M., 1991. Effect of chronic exposure of PCB (Aroclor 1254) on specific and non specific immune parameters in the rhesus (*Macaca mulatta*) monkey. *Fundamental and Applied Toxicology* 16, 772–786.
- Turyk, M., Anderson, H.A., Hanrahan, L.P., Falk, C., Steenport, D.N., Needham, L.L., Patterson Jr., D.G., Freels, S., Persky, V., Consortium, Great Lakes, 2006. Relationship of serum levels of individuals PCB, dioxin and furan congeners and DDE with Great Lakes sport-caught fish consumption. *Environmental Research* 100, 173–183.
- Ueno, D., Watanabe, M., Subramanian, A., Tanaka, H., Fillmann, G., Lam, P.K.S., Zheng, G.J., Muchtar, M., Razak, H., Prudente, M., 2005. Global pollution monitoring of polychlorinated dibenzo-*p*-dioxins (PCDDs), furans (PCDFs) and coplanar polychlorinated biphenyls (coplanar PCBs) using skipjack tuna as bioindicator. *Environmental Pollution* 136, 303–313.
- Vaclavik, E., Tjonneland, A., Stripp, C., Overvad, K., Weber, J.P., Raaschou-Nielsen, O., 2006. Organochlorines in Danish women: predictors of adipose tissue concentrations. *Environmental Research* 100, 362–370.
- Van den Berg, M., Birnbaum, L., Bosveld, A.T., Brunstrom, B., Cook, P., Feeley, M., Giesy, J.P., Hanberg, A., Hasegawa, R., Kennedy, S.W., Kubiak, T., Larsen, J.C., van Leeuwen, F.X., Liem, A.K., Nolt, C., Peterson, R.E., Poellinger, L., Safe, S., Schrenk, D., Tillitt, D., Tysklind, M., Younes, M., Waern, F., Zacharewski, T., 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environmental Health Perspective* 106, 775–792.
- Van den Berg, M., Birnbaum, L.S., Denison, M., De Vito, M., Farland, W., Feeley, M., Fiedler, H., Hakansson, H., Hanberg, A., Haws, L., Rose, M., Safe, S., Schrenk, D., Tohyama, C., Tritscher, A., Tuomisto, J., Tysklind, J.M., Walker, N., Richard, E., Peterson, R.E., 2005. The 2005 World Health Organization reevaluation of human and mammalian Toxic Equivalency Factors for dioxins and dioxin-like compounds. *Toxicological Sciences* 93, 223–241.
- Van Duursen, M.B., Sanderson, J.T., van der Bruggen, M., van der Linden, J., van den Berg, M., 2003. Effects of several dioxin-like compounds on estrogen metabolism in the malignant MCF-7 and nontumorigenic MCF-10 A human mammary epithelial cell lines. *Toxicology and Applied Pharmacology* 190, 241–250.
- Veza, C.M., Walker, N.J., Olson, J.R., 2004. Subchronic Exposure to TCDD, PeCDF, PCB126, and PCB153: Effect on Hepatic Gene Expression. *Environmental Health Perspective* 112, 1636–1644.
- Vives, I., Grimalt, J.O., Ventura, M., Catalan, J., Rossenland, B.O., 2005. Age dependence of the accumulation of organochlorine pollutants in brown trout (*Salmo trutta*) from a remote high mountain lake (Redó, Pyrenees). *Environmental Pollution* 133, 343–350.
- WHO Regional Office for Europe, 2000. Air Quality Guidelines, second ed. Copenhagen, Denmark. Cap 5.10 PCBs.
- Zeiger, M., Haag, R., Hacle, J., Schrenk, D., Schmitz, H.J., 2001. Inducing effects of dioxin like polychlorinated biphenyls on CYP1A in the human hepatoblastoma cell line HepG2, the rat hepatoma cell line H4IIE, and rat primary hepatocytes: Comparison of relative potencies. *Toxicological Sciences* 63 (1), 65–73.
- Zhou, W., Zhai, Z., Wang, Z., Wang, L., 2005. Estimation of *n*-octanol/water partition coefficients (*K*<sub>ow</sub>) of all PCB congeners by density functional theory. *Journal of Molecular Structure* 755, 137–145.